

Laboratorio Clinico Generale con aree specialistiche: Patologia Clinica e Sostanze d'Abuso - Microbiologia e Virologia

PUNTI PRELIEVI

Mod37E-rev4 del 19/04/2024 Inglese

URINE COLLECTION FOR CHLAMYDIA, URINE COLTURE, RANDOM URINE SAMPLE.

Equipment:

- Two Steril urine cups
- antiseptic soap and water
- antiseptic skin towelette, washcloth or sponge

You have to wait one hour after you last urinated before collecting a urine sample.

Female Collection

- 1. Wash hands thoroughly with soap and water. Dry with paper towel
- 2. With one hand, spread genital skin folds apart.
- 3. Using antiseptic skin towelette, wash vulva, wiping from front to back
- 4. Discard towelette
- 5. Repeat, wiping from front to back.
- 6. As you begin to urinate, collect the initial portion of your urine stream (first-catch) in the first container. Do not touch the inside of the container.
- 7. Stop collection when container is about half full . Screw cap on container
- 8. Void midstream portion into the second sterile container. Do not touch the inside of the container.
- Wash hands
- 10. Give samples to lab personnel. If there is a delay, specimen should be refrigerated.

Male Collection

- 1. Wash hands thoroughly with soap and water. Dry with paper towel.
- 2. Pull back foreskin, if uncircumcised and cleanse glans penis with antiseptic skin towelette.
- 3. As you begin to urinate, collect the initial portion of your urine stream (first-catch) in the first container. Do not touch the inside of the container.
- 4. Stop collection when container is about half full . Screw cap on container
- 5. Void midstream portion into the second sterile container. Do not touch the inside of the container.
- 6. Screw cap on container.
- 7. Wash hands.
- 8. Give samples to lab personnel. If there is a delay, specimen should be refrigerated.

La Direzione del Laboratorio



