

 <p>OSPEDALE SAN RAFFAELE</p>	<p>Laboratorio Clinico Generale con aree specialistiche: Patologia Clinica e Sostanze d'Abuso - Microbiologia e Virologia</p>	<p>PUNTI PRELIEVI ESTERNI Mod5E-rev4 del 19/02/2024 Inglese</p>
---	--	--

24-Hour Urine Test, Collection Instructions for 5-Hydroxyindoleacetic acid, Catecholamines, Vanillylmandelic acid VMA

The information your doctor receives from your test(s) is only as accurate as the collection procedure you follow. Therefore, it is important that you collect the urine in the following manner:

Before starting the urinary collection add 10 mL of **Hydrochloric Acid** given by the outpatient clinic.

1. You must begin all 24-hour urine collections with an empty bladder. When you wake up the morning of the day the collection is to begin, empty your bladder, discard the urine and note the time.
2. Collect all urine throughout the day and night. **Keep samples refrigerated at all times unless otherwise specified.**
3. The next morning wake up at the same time you did the day you began the collection, empty your bladder and add this to the urine you collected the day before. This completes the 24-hour urine collection.
4. Return specimen to the lab site as soon as possible after collection.

IMPORTANT: The bottle contains Hydrochloric Acid: DON'T DRINK, DO NOT LEAVE UNATTENDED. In case of ingestion, go to the Emergency hospital. Avoid contact with eyes and skin, otherwise wash carefully with a lot of water. Hazard Statements:

H314: Causes severe skin burns and eye damage



IMPORTANT: Before you stop taking any medications you must check with your physician!

The following is a list of preparation requirements for tests requiring 24 hour urine collection:

5-HIAA –Patients should avoid for a 72-hour period prior to and during collection the following foods and drugs or any compound containing these drugs; avocados, bananas, eggplant, plantain, plums, tomatoes, pineapples, walnuts, taking cough syrup containing guaifenesin, Tylenol, Empirin, Acetanilid, Glyceryl Guaiacolate, Mephenesin, Methocarbamol, Reserpine, Chlorpromazine, Promazine, Imipramine, Isoniazid, MAO Inhibitors, Methenamine, Methyldopa and Phenothiazines.

Catecholamines – Avoid patient stress. Many drugs (reserpine and alpha methyldopam, levadopa, monoamine oxidase inhibitors and sympathomimetic amines) may interfere and should be discontinued two weeks prior to specimen collection. Nose drops, sinus and cough medicines, bronchodilators and appetite suppressant– agonists, calcium channel blockers, converting enzyme - and β -blockers, and laetalol may interfere. Mandelamine® interferes, but thiazides do not. Caffeine products should be avoided before and during collection. The patient should not be subjected to hypoglycemia or exertion. Increased intracranial pressure and clonidine withdrawal can cause false-positive results.

Vanillylmandelic Acid VMA – Patients should avoid for 72 hours prior to collection salicylates, caffeine, phenothiazine and antihypertension agents. Also coffee, tea, chocolate, fruit (especially bananas and any vanilla containing substances).

La Direzione del Laboratorio

Ospedale San Raffaele S.r.l.
Istituto di Ricovero e Cura a Carattere Scientifico

Via Olgettina 60 – 20132 Milano (MI) | Tel. +39 02.26431 | info@hsr.it
C.F., P.IVA e Reg. Imp. Milano 07636600962 – C.C.I.A.A. 1972938
Capitale Sociale € 60.817.200 i.v.

www.hsr.it

Sistema Sanitario  Regione Lombardia

 **UniSR**
Università Vita-Salute
San Raffaele